

THE GRAM MATRIX OF A TEMPERLEY-LIEB ALGEBRA IS SIMILAR TO THE MATRIX OF CHROMATIC JOINS

QI CHEN AND JOZEF H. PRZYTICKI

Dedicated to the memory of Xiao-Song Lin (1957-2007)

To appear in Communications in Contemporary Mathematics

1. INTRODUCTION

Rodica Simion noticed experimentally that matrices of chromatic joins (introduced by W. Tutte in [Tu2]) and the Gram matrix of the Temperley-Lieb algebra, have the same determinant, up to renormalization. In the type A case, she was able to prove this by comparing the known formulas: by Tutte and R. Dahab [Tu2, Dah], in the case of chromatic joins, and by P. Di Francesco, and B. Westbury [DiF, We] (based on the work by K. H. Ko and L. Smolinsky [KS]) in the Temperley-Lieb case; see [CSS]. She then asked for a direct proof of this fact [CSS], [Sch], Problem 7.

The type B analogue was an open problem central to the work of Simion [Sch]. She demonstrated strong evidence that the type B Gram determinant of the Temperley-Lieb algebra is equal to the determinant of the matrix of type B chromatic joins, after a substitution similar to that in type A , cf. [Sch].

In this paper we show that the matrix J_n of chromatic joins and the Gram matrix G_n of the Temperley-Lieb algebra are similar (after rescaling), with the change of basis given by diagonal matrices. More precisely we prove the following two results:

Theorem A. *We have $J_n^A(\delta^2) = PG_n^A(\delta)P$, where $P = (p_{ij})$ is a diagonal matrix with $p_{ii}(\delta) = \delta^{\text{bk}(\pi_i)-n/2}$; here $\text{bk}(\pi_i)$ denotes the number of blocks in the type A non-crossing n -partition π_i ; see 2.1 of Section 2 for precise definitions.*

Theorem B. *We have $J_n^B(\delta^2) = P^B G_n^B(1, \delta) P^B$ where $P^B = (p_{ij}^B)$ is a diagonal matrix with $p_{ii}^B(\delta) = \delta^{\text{nzbk}(\pi_i)-n/2}$; here $\text{nzbk}(\pi_i)$ denotes half of the number of non-zero blocks in the type B non-crossing n -partition π_i ; see 2.2 of Section 2 for precise definitions.*

2. DEFINITIONS AND NOTATION

In the description of the matrix of chromatic joins (of type A and B) we follow V. Reiner, R. Simion and F. Schmidt [Rei, Sim, Sch].

2.1. The type A case. An n -partition of type A is a partition π of the n element set $\{1, 2, \dots, n\}$ into blocks. The number of blocks is denoted by $\text{bk}(\pi)$. To represent π pictorially, we place the numbers $1, 2, \dots, n$ anti-clockwise around the boundary circle of the unit disk and draw a chord, called a *connection chord*, in the disk between two numbers $i < j$ if they are in the same block of π and there is no k in the same block with $i < k < j$. We say that π is *non-crossing* if all connection chords can be drawn without crossing each other. Notice that each block is represented by a tree. Denote the set of all non-crossing n -partitions of type A by Π_n^A . On the other hand if $n = 2m$ is even, we have *bipartitions* of $2m$ points of type A , those $2m$ -partitions of type A with every block containing exactly 2 numbers. Denote the set of all non-crossing $2m$ -bipartitions of type A by Γ_m^A . We have a bijection $\varphi_A : \Pi_n^A \rightarrow \Gamma_n^A$ realized by considering the boundary arcs of a regular neighborhood of the connection chords (see Fig. 1 and compare Fig. 2, Fig. 3).




FIGURE 1. The bijection $\varphi_A : \Pi_3^A \rightarrow \Gamma_3^A$.

2.2. The type B case. An n -partition of type B is a partition π of the $2n$ element set $\{+1, +2, \dots, +n, -1, -2, \dots, -n\}$ into blocks with the property that for any block K of π , its opposite $-K$ is also a block of π , and that there is at most one invariant block (called the zero block[†]) for which $K = -K$. Since all non-zero blocks occur in pairs $\pm K$ one defines $\text{nzbk}(\pi)$ as half of the number of all non-zero blocks. To represent π pictorially, we place the numbers $+1, +2, \dots, +n, -1, -2, \dots, -n$ anti-clockwise around the boundary circle of a disk and draw a connection chord in the disk between two numbers $i < j^{\ddagger}$ if they are in the same block of π and there is no k in the same block with $i < k < j$. Then π is said to be *non-crossing* if all connection chords can be drawn without crossing each other.[§] Denote the set of all non-crossing n -partitions of type B by Π_n^B . The set Π_n^B is illustrated in Fig. 2.

[†]For topologists the term invariant block is more natural than zero block so we use this names interchangeably in the paper.

[‡]Here we use the order $+1 < +2 < \dots < n < -1 < -2 < \dots < -n$.

[§]The non-crossing condition forces a partition to have at most one zero block.

On the other hand if $n = 2m$ is even, we have bipartitions of $4m$ points of type




FIGURE 2. The pictorial representation of Π_2^B .

B , those $2m$ -partitions of type B with every block containing exactly 2 numbers. Denote the set of all non-crossing $2m$ -bipartitions of type B by Γ_m^B . Similar to the type A case we have a bijection $\varphi_B : \Pi_n^B \rightarrow \Gamma_n^B$ realized by considering the boundary arcs of a regular neighborhood of the connection chords (see Fig. 3).




FIGURE 3. The bijection $\varphi_B : \Pi_2^B \rightarrow \Gamma_2^B$.

2.3. The matrices. For any n -partitions π and π' of type A (resp. B), denote by $\pi \vee \pi'$ the finest n -partition (not necessarily non-crossing) of type A (resp. B) that is coarser than both π and π' . The *matrix of chromatic joins* of type A and B are respectively:

$$(J_n^A(\delta))_{\pi, \pi' \in \Pi_n^A} = \delta^{\text{bk}(\pi \vee \pi')} \quad \text{and} \quad (J_n^B(\delta))_{\pi, \pi' \in \Pi_n^B} = \delta^{\text{nzbk}(\pi \vee \pi')}.$$

For any $2n$ -bipartitions π and π' of type A (resp. B), one can glue them along the boundary circles respecting the labels. The result, denoted $\pi \vee \pi'$, is a collection of disjoint circles on a 2-sphere. The *Gram matrix of Temperley-Lieb algebra* of type A and B are respectively[¶]:

$$(G_n^A(\delta))_{\pi, \pi' \in \Gamma_n^A} = \delta^{c(\pi \vee \pi')} \quad \text{and} \quad (G_n^B(\alpha, \delta))_{\pi, \pi' \in \Gamma_n^B} = \alpha^{c_0(\pi \vee \pi')} \delta^{c_d(\pi \vee \pi')},$$

where $c(\pi \vee \pi')$ is the number of circles, $c_0(\pi \vee \pi')$ is the number of zero (i.e. invariant) circles C with $C = -C$, and $c_d(\pi \vee \pi')$ is the number of pairs of non-zero circles $C, -C$ with $C \neq -C$ in $\pi \vee \pi'$.

[¶]The matrix $G_n^A(\delta)$ was first used by H. Morton and P. Traczyk to find a basis of the Kauffman bracket skein module of a tangle [MT], and played an important role in Lickorish's approach to Witten-Reshetikhin-Turaev invariants of 3-manifolds [Li]. The matrix $G_n^B(1, \delta)$ was first considered by Rodica Simion in 1998; compare [Sch].

3. PROOF OF THEOREMS A AND B

Proof of Theorem A. For $\pi_i \in \Pi_n^A$ let $b_i := \varphi_A(\pi_i) \in \Gamma_n^A$. Since $c(b_i \vee b_j)$ is also equal to the number of boundary components of the regular neighborhood of the pictorial representation of $\pi_i \vee \pi_j$ we have $2 \text{bk}(\pi_i \vee \pi_j) = c(b_i \vee b_j) + \text{bk}(\pi_i) + \text{bk}(\pi_j) - n = \text{bk}(\pi_i) - \frac{n}{2} + c(b_i \vee b_j) + \text{bk}(\pi_j) - \frac{n}{2}$. The formula can be obtained from the expression for the Euler characteristic of a plane graph: Let G_π be a graph corresponding to the non-crossing partition π . G_π is a forest of n vertices and $n - \text{bk}(\pi)$ edges. Similarly, let $G_{\pi_i \vee \pi_j}$ be the graph corresponding to $\pi_i \vee \pi_j$. We should stress that $\pi_i \vee \pi_j$ does not have to be a noncrossing partition and that the graph $G_{\pi_i \vee \pi_j}$ is a plane graph obtained by putting G_{π_i} inside a disk and G_{π_j} outside the disk with $G_{\pi_i} \cap G_{\pi_j}$ composed of the n points on the unit circle

(e.g.:   , or  ).

By construction, $G_{\pi_i \vee \pi_j}$ is a plane graph of n vertices and $\text{bk}(\pi_i \vee \pi_j)$ components. It has $E(G_{\pi_i \vee \pi_j}) = E(G_{\pi_i}) + E(G_{\pi_j}) = n - \text{bk}(\pi_i) + n - \text{bk}(\pi_j)$ edges. Furthermore, if we embed $G_{\pi_i \vee \pi_j}$ in a disjoint union of $\text{bk}(\pi_i \vee \pi_j)$ 2-spheres (each component of $G_{\pi_i \vee \pi_j}$ in a different sphere) we can identify $c(b_i \vee b_j)$ with the number of regions of the embedded graph. The Euler characteristic is on the one hand equal to $2\text{bk}(\pi_i \vee \pi_j)$ and on the other hand equal to $n - E(G_{\pi_i \vee \pi_j}) + c(b_i \vee b_j) = c(b_i \vee b_j) + \text{bk}(\pi_i) + \text{bk}(\pi_j) - n$, as needed.

Theorem A follows directly from the formula. $\square$

Proof of Theorem B. For $\pi_i \in \Pi_n^B$ let $b_i := \varphi_B(\pi_i) \in \Gamma_n^B$ and $\text{bk}_0(\pi_i)$ be the number of zero blocks of π_i . Recall that $c_0(b_i \vee b_j)$ is the number of zero (i.e. invariant) components of $b_i \vee b_j$. As in the type A case we have $2 \text{bk}(\pi_i \vee \pi_j) = c(b_i \vee b_j) + \text{bk}(\pi_i) + \text{bk}(\pi_j) - 2n$. Furthermore, we have (see Lemma 1):

$$2 \text{bk}_0(\pi_i \vee \pi_j) = c_0(b_i \vee b_j) + \text{bk}_0(\pi_i) + \text{bk}_0(\pi_j).$$

(Notice that $\text{bk}_0(\pi_i \vee \pi_j)$ can be equal to 2, 1, or 0.) From these we conclude that: $2 \text{nzbk}(\pi_i \vee \pi_j) = c_d(b_i \vee b_j) + \text{nzbk}(\pi_i) + \text{nzbk}(\pi_j) - n$. Thus Theorem B follows. $\square$

Lemma 1. *The zero blocks and zero components satisfy the following identity:*

$$2 \text{bk}_0(\pi_i \vee \pi_j) = c_0(b_i \vee b_j) + \text{bk}_0(\pi_i) + \text{bk}_0(\pi_j),$$

where $b_i = \varphi_B(\pi_i)$ and $b_j = \varphi_B(\pi_j)$.

Proof. The lemma reflects the basic properties of a 2-sphere with an involution fixing two points and its compact invariant submanifolds.

To demonstrate the formula we consider all cases of blocks of π_i , π_j and $\pi_i \vee \pi_j$ divided into four classes:

(1) If K is a non-zero (i.e. non-invariant) block of $\pi_i \vee \pi_j$ then all its constituent blocks in π_i and π_j are non-zero blocks and the boundary components of a regular neighborhood of the geometric realization of K (denoted ∂K), that is circles in $b_i \vee b_j$ corresponding to K , are non-invariant (non-zero) curves, i.e. not in $c_0(b_i \vee b_j)$.

- (2) If K is a zero block of $\pi_i \vee \pi_j$ but all its constituent blocks in π_i and π_j are non-zero blocks, then exactly two components of ∂K are invariant curves.
- (3) If K is a zero block of $\pi_i \vee \pi_j$ and exactly one constituent block is a zero block then exactly one component of ∂K is an invariant curve.
- (4) If K is a zero block of $\pi_i \vee \pi_j$ and exactly two constituent blocks are zero-blocks (necessarily one in π_i and one in π_j) then no component of ∂K is an invariant curve.

These conditions taken together prove the formula in Lemma 1. $\square$

4. COROLLARIES

Theorem B and the results of [MS, CP] allow us to answer Problems 1 and 2 of [Sch] about a formula for the determinant of the type- B matrix of chromatic joins:

Corollary 2.

$$\det(J_n^B(\delta^2)) = \prod_{i=1}^n (T_i(\delta)^2 - 1)^{\binom{2n}{n-i}}$$

where $T_i(\delta)$ is the Chebyshev polynomial of the first kind:

$$T_0 = 2, \quad T_1 = \delta, \quad T_i = \delta T_{i-1} - T_{i-2}.$$

The matrix $J_n^B(\delta)$ can be generalized to a matrix of two variables as follows:

$$(J_n^B(\alpha, \delta))_{\pi, \pi' \in \Pi_n^B} = \alpha^{\text{bk}_0(\pi \vee \pi')} \delta^{\text{nzbk}(\pi \vee \pi')}.$$

It follows from Lemma 1 that

$$J_n^B(\alpha^2, \delta^2) = P_n^B(\alpha, \delta) G_n^B(\alpha, \delta) P_n^B(\alpha, \delta),$$

where $P_n^B(\alpha, \delta) = (p_{ij})$ is a diagonal matrix with $p_{ii}(\alpha, \delta) = \alpha^{\text{bk}_0(\pi_i)} \delta^{\text{nzbk}(\pi_i) - n/2}$. Furthermore, $\det P_n^B(\alpha, \delta) = \alpha^{\frac{1}{2} \binom{2n}{n} **}$ and thus we have:

Corollary 3.

$$\det J_n^B(\alpha^2, \delta^2) = \alpha^{\binom{2n}{n}} G_n^B(\alpha, \delta) = \alpha^{\binom{2n}{n}} \prod_{i=1}^n (T_i(\delta)^2 - \alpha^2)^{\binom{2n}{n-i}}.$$

Remark 4. Consider the Gram matrix of type B based on non-crossing connections in an annulus.^{††} This matrix is the same as the one considered before in Theorem B via the branched cover described in Fig. 4.

^{**}This follows from Proposition 3 of [Rei], which asserts that there exists a fixed-point free involution γ on Π_n^B such that $\text{bk}_0(\pi) + \text{bk}_0(\gamma(\pi)) = 1$ and $\text{nzbk}(\pi) + \text{nzbk}(\gamma(\pi)) = n$.

^{††}This interpretation of the Gram matrix of type B Temperley-Lieb algebra is mentioned in [Sch] as an annular skein matrix and utilized in [MS] and [CP].




FIGURE 4. The double branch cover $pr : D^2 \rightarrow D^2$ with the “cutting” arc S

REFERENCES

- [CP] Q. Chen, J. H. Przytycki, The Gram determinant of the type B Temperley-Lieb algebra, e-print: <http://arxiv.org/abs/0802.1083>
- [CSS] A. Copeland, F. Schmidt, R. Simion, Note on two determinants with interesting factorizations, *Discrete Mathematics*, 256:449–458, 2002.
- [Dah] R. Dahab, The Birkhoff-Lewis equation, PhD dissertation, University of Waterloo, 1993.
- [DiF] P. Di Francesco, Meander determinants, *Comm. Math. Phys.*, 191:543–583, 1998.
- [KS] K. H. Ko, L. Smolinsky, A combinatorial matrix in 3-manifold theory, *Pacific Journ. Math.*, 149(2), 1991, 319–336.
- [Li] W. B. R. Lickorish, Invariants for 3-manifolds from the combinatorics of the Jones polynomial, *Pacific Journ. Math.*, 149(2):337–347, 1991.
- [MS] P. P. Martin, H. Saleur, On an Algebraic Approach to Higher Dimensional Statistical Mechanics, *Commun. Math. Phys.* 158, 1993, 155–190; e-print: <http://front.math.ucdavis.edu/9208.3061>
- [MT] H. R. Morton, P. Traczyk, Knots and algebras, *Contribuciones Matematicas en homenaje al profesor D. Antonio Plans Sanz de Bremond*, ed. E. Martin-Peinador and A. Rodez Usan, University of Zaragoza, pp. 201–220, 1990.
- [Rei] V. Reiner, Non-crossing partitions for classical reflection groups. *Discrete Math.*, 177:(195–222), 1997.
- [Sch] F. Schmidt, Problems related to type-A and type-B matrices of chromatic joins, *Advances in Applied Mathematics*, 32:(380–390), 2004.
- [Sim] R. Simion, Noncrossing partitions, *Discrete Math.*, 217:367–409, 2000.
- [Tu2] W. T. Tutte, The matrix of chromatic joins, *J. Combin. Theory Ser. B*, 57:(269–288), 1993.
- [We] B. W. Westbury, The representation theory of the Temperley-Lieb algebras. *Math. Z.*, 219(4):539–565, 1995.

Department of Mathematics
Winston-Salem State University
Winston Salem, NC 27110, USA
chenqi@wssu.edu

Department of Mathematics
The George Washington University
Washington, DC 20052, USA
przytyck@gwu.edu